

# UNITED STATES

## TWELFTH GRADE DRUG USE

	2002	2001	2000	1999	1998	1997	1996	1995	1994	1993	1992
<b>Percent Currently Using Drug</b>											
Alcohol	48.6	49.8	50.0	51.0	52.0	52.7	50.8	51.3	50.1	48.6	
Been Drunk Within Past 30 days	30.3	32.7	32.3	32.9	32.9	34.2	31.3	33.2	30.8	28.9	29.9
Binge Alcohol Use *	28.6	29.7	30.0	30.8	31.5	31.3	30.2	29.8	28.2	27.5	27.9
Cigarettes	26.7	29.5	31.4	34.6	35.1	36.5	34.0	33.5	31.2	29.9	27.8
Smokeless Tobacco	6.5	7.8	7.6	8.4	8.8	9.7	9.8	12.2	11.1	10.7	11.4
Any Illicit Drug	25.4	25.7	24.9	25.9	25.6	26.2	24.6	23.8	21.9	18.3	14.4
Marijuana/Hashish	21.5	22.4	21.6	23.1	22.8	23.7	21.9	21.2	19.0	15.5	11.9
Inhalants	1.5	1.7	2.2	2.0	2.3	2.5	2.5	3.2	2.7	2.5	2.3
Hallucinogens	2.3	3.3	2.6	3.5	3.8	3.9	3.5	4.4	3.1	2.7	2.1
LSD	0.7	2.3	1.6	2.7	3.2	3.1	2.5	4.0	2.6	2.4	2.0
PCP	0.4	0.5	0.9	0.8	1.0	0.7	1.3	0.6	0.7	1.0	0.6
Cocaine (all forms)	2.3	2.1	2.1	2.6	2.4	2.3	2.0	1.8	1.5	1.3	1.3
Crack Cocaine	1.2	1.1	1.0	1.1	1.0	0.9	1.0	1.0	0.8	0.7	0.6
Heroin	0.5	0.4	0.7	0.5	0.5	0.5	0.5	0.6	0.3	0.2	0.3
Other Narcotics	3.1	3.0	2.9	2.6	2.4	2.3	2.0	1.8	1.5	1.3	1.2
Amphetamines	5.5	5.6	5.0	4.5	4.6	4.8	4.1	4.0	4.0	3.7	2.8
Methamphetamine	1.7	1.5	1.9	1.7							
Crystal Methamphetamine (Ice)	1.2	1.1	1.0	0.8	1.2	0.8	1.1	1.1	0.7	0.6	0.5
Sedatives								2.3	1.8	1.3	1.2
Barbiturates	3.2	2.8	3.0	2.6	2.6	2.1	2.1	2.2	1.7	1.3	1.1
Methaqualone								0.4	0.4	0.1	0.4
Tranquilizers	3.3	2.9	2.6	2.5	2.4	1.8	2.0	1.8	1.4	1.2	1.0
Steroids	1.4	1.3	0.8	0.9	1.1	1.0	0.7	0.7	0.9	0.7	0.6
MDMA (Ecstasy)	2.4	2.8	3.6	2.5	1.5	1.6	2.0				

	2002	2001	2000	1999	1998	1997	1996	1995	1994	1993	1992
<b>Percent Ever Using Drug</b>											
Alcohol	78.4	79.7	80.3	80.0	81.4	81.7	79.2	80.7	80.4	80.0	
Ever Been Drunk	61.6	63.9	62.3	62.3	62.4	64.2	61.8	63.2	62.9	62.5	63.4
Cigarettes	57.2	61.0	62.5	64.6	65.3	65.4	63.5	64.2	62.0	61.9	61.8
Smokeless Tobacco	18.3	19.7	23.1	23.4	26.2	25.3	29.8	30.9	30.7	31.0	32.4
Any Illicit Drug	53.0	53.9	54.0	54.7	54.1	54.3	50.8	48.4	45.6	42.9	40.7
Marijuana/Hashish	47.8	49.0	48.8	49.7	49.1	49.6	44.9	41.7	38.2	35.3	32.6
Inhalants	11.7	13.0	14.2	15.4	15.2	16.1	16.6	17.4	17.7	17.4	16.6
Hallucinogens	12.0	14.7	13.0	13.7	14.1	15.1	14.0	12.7	11.4	10.9	9.2
LSD	8.4	10.9	11.1	12.2	12.6	13.6	12.6	11.7	10.5	10.3	8.6
PCP	3.1	3.5	3.4	3.4	3.9	3.9	4.0	2.7	2.8	2.9	2.4
Cocaine (all forms)	7.8	8.2	8.6	9.8	9.3	8.7	7.1	6.0	5.9	6.1	6.1
Crack Cocaine	3.8	3.7	3.9	4.6	4.4	3.9	3.3	3.0	3.0	2.6	2.6
Heroin	1.7	1.8	2.4	2.0	2.0	2.1	1.8	1.6	1.2	1.1	1.2
Other Narcotics	10.1	9.9	10.6	10.2	9.8	9.7	8.2	7.2	6.6	6.4	6.1
Amphetamines	16.8	16.2	15.6	16.3	16.4	16.5	15.3	15.3	15.7	15.1	13.9
Methamphetamine	6.7	6.9	7.9	8.2							
Crystal Methamphetamine (Ice)	4.7	4.1	4.0	4.8	5.3	4.4	4.4	3.9	3.4	3.1	2.9
Sedatives								7.6	7.3	6.4	6.1
Barbiturates	9.5	8.7	9.2	8.9	8.7	8.1	7.6	7.4	7.0	6.3	5.5
Methaqualone								1.2	1.4	0.8	1.6
Tranquilizers	11.4	10.3	8.9	9.3	8.5	7.8	7.2	7.1	6.6	6.4	6.0
Steroids	4.0	3.7	2.5	2.9	2.7	2.4	1.9	2.3	2.4	2.0	2.1
MDMA (Ecstasy)	10.5	11.7	11.0	8.0	5.8	6.9	6.1				

=> *Italicized* percentages in 2001 data are different than previously reported due to revision of estimates by the Monitoring the Future program

\* "Binge" alcohol use in the Monitoring the Future Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past two weeks

Source: [71]